

# Introduction to Mobile Robotics

## Compact Course on Linear Algebra


Wolfram Burgard, Maren Bennewitz,  
Diego Tipaldi, Luciano Spinello



# Vectors


- Arrays of numbers
- Vectors represent a point in a  $n$  dimensional space

$$(a_1) \quad \begin{pmatrix} a_1 \\ a_2 \end{pmatrix} \quad \begin{pmatrix} a_1 \\ a_2 \\ \vdots \\ a_n \end{pmatrix}$$



# Vectors: Scalar Product

- Scalar-Vector Product  $ka$
- Changes the length of the vector, but **not** its direction




# Vectors: Sum

- Sum of vectors (is commutative)

$$\begin{pmatrix} a_1 \\ a_2 \\ \vdots \\ a_n \end{pmatrix} + \begin{pmatrix} b_1 \\ b_2 \\ \vdots \\ b_n \end{pmatrix} = \begin{pmatrix} b_1 \\ b_2 \\ \vdots \\ b_n \end{pmatrix} + \begin{pmatrix} a_1 \\ a_2 \\ \vdots \\ a_n \end{pmatrix}$$

- Can be visualized as “chaining” the vectors.




# Vectors: Dot Product

- Inner product of vectors (is a scalar)

$$\mathbf{a} \cdot \mathbf{b} = \mathbf{b} \cdot \mathbf{a} = \sum_i a_i b_i$$


- If one of the two vectors, e.g.  $\mathbf{a}$ , has  $\|\mathbf{a}\| = 1$  the inner product  $\mathbf{a} \cdot \mathbf{b}$  returns the length of the projection of  $\mathbf{b}$  along the direction of  $\mathbf{a}$



- If  $\mathbf{a} \cdot \mathbf{b} = 0$ , the two vectors are **orthogonal**


# Vectors: Linear (In)Dependence

- A vector  $\mathbf{b}$  is **linearly dependent** from  $\{\mathbf{a}_1, \mathbf{a}_2, \dots, \mathbf{a}_n\}$  if  $\mathbf{b} = \sum_i k_i \mathbf{a}_i$
- In other words, if  $\mathbf{b}$  can be obtained by summing up the  $\mathbf{a}_i$  properly scaled
- If there exist no  $\{k_i\}$  such that  $\mathbf{b} = \sum_i k_i \mathbf{a}_i$  then  $\mathbf{b}$  is independent from  $\{\mathbf{a}_i\}$



# Vectors: Linear (In)Dependence

- A vector  $\mathbf{b}$  is **linearly dependent** from  $\{\mathbf{a}_1, \mathbf{a}_2, \dots, \mathbf{a}_n\}$  if  $\mathbf{b} = \sum_i k_i \mathbf{a}_i$
- In other words, if  $\mathbf{b}$  can be obtained by summing up the  $\mathbf{a}_i$  properly scaled
- If there exist no  $\{k_i\}$  such that  $\mathbf{b} = \sum_i k_i \mathbf{a}_i$  then  $\mathbf{b}$  is independent from  $\{\mathbf{a}_i\}$



# Matrices

- A matrix is written as a table of values

$$\mathbf{A} = \begin{pmatrix} a_{11} & a_{12} & \cdots & a_{1m} \\ a_{21} & a_{22} & \cdots & a_{2m} \\ \vdots & & & \\ a_{n1} & a_{n2} & \cdots & a_{nm} \end{pmatrix}$$

$$\mathbf{A} : n \times m$$

↑            ↑  
rows        columns

- **1<sup>st</sup> index** refers to the **row**
- **2<sup>nd</sup> index** refers to the **column**
- Note: a d-dimensional vector is equivalent to a  $d \times 1$  matrix



# Matrices as Collections of Vectors

- Column vectors

$$\mathbf{A} = \begin{pmatrix} \boxed{\begin{matrix} a_{11} \\ a_{21} \\ \vdots \\ a_{n1} \end{matrix}} & \boxed{\begin{matrix} a_{12} \\ a_{22} \\ \vdots \\ a_{n2} \end{matrix}} & \cdots & \boxed{\begin{matrix} a_{1m} \\ a_{2m} \\ \vdots \\ a_{nm} \end{matrix}} \end{pmatrix}$$

$\begin{matrix} \uparrow & \uparrow & & \uparrow \\ \mathbf{a}_{*1} & \mathbf{a}_{*2} & \cdots & \mathbf{a}_{*m} \end{matrix}$

# Matrices as Collections of Vectors

- Row vectors

$$\mathbf{A} = \begin{pmatrix} \boxed{a_{11} \quad a_{12} \quad \cdots \quad a_{1m}} \\ \boxed{a_{21} \quad a_{22} \quad \cdots \quad a_{2m}} \\ \vdots \\ \boxed{a_{n1} \quad a_{n2} \quad \cdots \quad a_{nm}} \end{pmatrix} \begin{matrix} \longrightarrow \\ \longrightarrow \\ \longrightarrow \\ \longrightarrow \end{matrix} \begin{pmatrix} \mathbf{a}_{1*}^T \\ \mathbf{a}_{2*}^T \\ \vdots \\ \mathbf{a}_{n*}^T \end{pmatrix}$$

# Important Matrices Operations

- Multiplication by a scalar
- Sum (commutative, associative)
- Multiplication by a vector
- Product (not commutative)
- Inversion (square, full rank)
- Transposition

# Scalar Multiplication & Sum

- In the scalar multiplication, every element of the vector or matrix is multiplied with the scalar
- The sum of two vectors is a vector consisting of the pair-wise sums of the individual entries
- The sum of two matrices is a matrix consisting of the pair-wise sums of the individual entries

# Matrix Vector Product


- The  $j^{\text{th}}$  component of  $\mathbf{A}\mathbf{b}$  is the dot product  $\mathbf{a}_{j*}^T \cdot \mathbf{b}$ .
- The vector  $\mathbf{A}\mathbf{b}$  is linearly dependent from the column vectors  $\{\mathbf{a}_{*i}\}$  with coefficients  $\{b_i\}$

$$\mathbf{A}\mathbf{b} = \begin{pmatrix} \mathbf{a}_{1*}^T \\ \mathbf{a}_{2*}^T \\ \vdots \\ \mathbf{a}_{n*}^T \end{pmatrix} \cdot \mathbf{b} = \begin{pmatrix} \mathbf{a}_{1*}^T \cdot \mathbf{b} \\ \mathbf{a}_{2*}^T \cdot \mathbf{b} \\ \vdots \\ \mathbf{a}_{n*}^T \cdot \mathbf{b} \end{pmatrix} = \sum_k \mathbf{a}_{*k} b_k$$

row vectors                      column vectors

# Matrix Vector Product

- If the column vectors of  $\mathbf{A}$  represent a reference system, the product  $\mathbf{A}\mathbf{b}$  computes the global transformation of the vector  $\mathbf{b}$  according to  $\{\mathbf{a}_{*i}\}$



# Matrix Matrix Product

- Can be defined through
  - the dot product of row and column vectors
  - the linear combination of the columns of  $\mathbf{A}$  scaled by the coefficients of the columns of  $\mathbf{B}$


$$\begin{aligned}\mathbf{C} &= \mathbf{AB} \\ &= \begin{pmatrix} \mathbf{a}_{1*}^T \cdot \mathbf{b}_{*1} & \mathbf{a}_{1*}^T \cdot \mathbf{b}_{*2} & \cdots & \mathbf{a}_{1*}^T \cdot \mathbf{b}_{*m} \\ \mathbf{a}_{2*}^T \cdot \mathbf{b}_{*1} & \mathbf{a}_{2*}^T \cdot \mathbf{b}_{*2} & \cdots & \mathbf{a}_{2*}^T \cdot \mathbf{b}_{*m} \\ \vdots & & & \\ \mathbf{a}_{n*}^T \cdot \mathbf{b}_{*1} & \mathbf{a}_{n*}^T \cdot \mathbf{b}_{*2} & \cdots & \mathbf{a}_{n*}^T \cdot \mathbf{b}_{*m} \end{pmatrix} \\ &= \left( \mathbf{Ab}_{*1} \quad \mathbf{Ab}_{*2} \quad \cdots \quad \mathbf{Ab}_{*m} \right)\end{aligned}$$

 column vectors

# Matrix Matrix Product

- If we consider the second interpretation, we see that the columns of  $\mathbf{C}$  are the “transformations” of the columns of  $\mathbf{B}$  through  $\mathbf{A}$
- All the interpretations made for the matrix vector product hold

$$\begin{aligned}\mathbf{C} &= \mathbf{A}\mathbf{B} \\ &= \left( \mathbf{A}\mathbf{b}_{*1} \quad \mathbf{A}\mathbf{b}_{*2} \quad \dots \quad \mathbf{A}\mathbf{b}_{*m} \right) \\ \mathbf{c}_{*i} &= \mathbf{A}\mathbf{b}_{*i}\end{aligned}$$



column vectors



# Rank

- **Maximum** number of linearly independent rows (columns)
- Dimension of the **image** of the transformation  $f(\mathbf{x}) = A\mathbf{x}$
- When  $A$  is  $m \times n$  we have
  - $\text{rank}(A) \geq 0$  and the equality holds iff  $A$  is the null matrix
  - $\text{rank}(A) \leq \min(m, n)$
- Computation of the rank is done by
  - Gaussian elimination on the matrix
  - Counting the number of non-zero rows

# Inverse

$$\mathbf{AB} = \mathbf{I}$$

- If  $\mathbf{A}$  is a square matrix of full rank, then there is a unique matrix  $\mathbf{B}=\mathbf{A}^{-1}$  such that  $\mathbf{AB}=\mathbf{I}$  holds
- The  $i^{\text{th}}$  row of  $\mathbf{A}$  is and the  $j^{\text{th}}$  column of  $\mathbf{A}^{-1}$  are:
  - orthogonal (if  $i \neq j$ )
  - or their dot product is 1 (if  $i = j$ )

# Matrix Inversion

$$\mathbf{A}\mathbf{B} = \mathbf{I}$$

- The  $i^{\text{th}}$  column of  $\mathbf{A}^{-1}$  can be found by solving the following linear system:

$$\mathbf{A}\mathbf{a}^{-1}_{*i} = \mathbf{i}_{*i} \longleftarrow \text{This is the } i^{\text{th}} \text{ column of the identity matrix}$$

# Determinant (det)

- Only defined for **square matrices**
- The inverse of  $\mathbf{A}$  exists if and only if  $\det(\mathbf{A}) \neq 0$
- For  $2 \times 2$  matrices:

Let  $\mathbf{A} = [a_{ij}]$  and  $|\mathbf{A}| = \det(\mathbf{A})$ , then

$$\begin{vmatrix} a_{11} & a_{12} \\ a_{21} & a_{22} \end{vmatrix} = a_{11} \cdot a_{22} - a_{12} \cdot a_{21}$$

- For  $3 \times 3$  matrices the Sarrus rule holds:

$$\begin{vmatrix} a_{11} & a_{12} & a_{13} \\ a_{21} & a_{22} & a_{23} \\ a_{31} & a_{32} & a_{33} \end{vmatrix} = a_{11}a_{22}a_{33} + a_{12}a_{23}a_{31} + a_{13}a_{21}a_{32} \\ - a_{11}a_{23}a_{32} - a_{12}a_{21}a_{33} - a_{13}a_{22}a_{31}$$

# Determinant

- For **general**  $n \times n$  matrices?

Let  $\mathbf{A}_{ij}$  be the submatrix obtained from  $\mathbf{A}$  by deleting the  $i$ -th row and the  $j$ -th column

$$\begin{bmatrix} 1 & 2 & 5 & 0 \\ 2 & 3 & 4 & -1 \\ -5 & 8 & 0 & 0 \\ 0 & 4 & -2 & 0 \end{bmatrix} \quad \rightarrow \quad \mathbf{A}_{32} = \begin{bmatrix} 1 & 5 & 0 \\ 2 & 4 & -1 \\ 0 & -2 & 0 \end{bmatrix}$$

Rewrite determinant for  $3 \times 3$  matrices:

$$\begin{aligned} \det(\mathbf{A}^{3 \times 3}) &= a_{11}a_{22}a_{33} + a_{12}a_{23}a_{31} + a_{13}a_{21}a_{32} \\ &\quad - a_{11}a_{23}a_{32} - a_{12}a_{21}a_{33} - a_{13}a_{22}a_{11} \\ &= a_{11} \cdot \det(\mathbf{A}_{11}) - a_{12} \cdot \det(\mathbf{A}_{12}) + a_{13} \cdot \det(\mathbf{A}_{13}) \end{aligned}$$

# Determinant

- For **general**  $n \times n$  matrices?

$$\begin{aligned} \det(\mathbf{A}) &= a_{11}\det(\mathbf{A}_{11}) - a_{12}\det(\mathbf{A}_{12}) + \dots + (-1)^{1+n}a_{1n}\det(\mathbf{A}_{1n}) \\ &= \sum_{j=1}^n (-1)^{1+j}a_{1j}\det(\mathbf{A}_{1j}) \end{aligned}$$

Let  $\mathbf{C}_{ij} = (-1)^{i+j}\det(\mathbf{A}_{ij})$  be the  $(i,j)$ -cofactor, then

$$\begin{aligned} \det(\mathbf{A}) &= a_{11}\mathbf{C}_{11} + a_{12}\mathbf{C}_{12} + \dots + a_{1n}\mathbf{C}_{1n} \\ &= \sum_{j=1}^n a_{1j}\mathbf{C}_{1j} \end{aligned}$$

This is called the **cofactor expansion** across the first row

# Determinant

- **Problem:** Take a 25 x 25 matrix (which is considered small). The cofactor expansion method requires  $n!$  multiplications. For  $n = 25$ , this is  $1.5 \times 10^{25}$  multiplications for which a today supercomputer would take **500,000 years**.
- There are **much faster methods**, namely using **Gauss elimination** to bring the matrix into triangular form.

$$\mathbf{A} = \begin{bmatrix} d_1 & * & * & * \\ 0 & d_2 & * & * \\ 0 & 0 & d_3 & * \\ 0 & 0 & 0 & d_4 \end{bmatrix} \quad \det(\mathbf{A}) = \prod_{i=1}^n d_i$$

Because for **triangular matrices** the determinant is the product of diagonal elements


# Determinant: Properties

- **Row operations** ( $\mathbf{A}$  is still a  $n \times n$  square matrix)
  - If  $\mathbf{B}$  results from  $\mathbf{A}$  by interchanging two rows, then  $\det(\mathbf{B}) = -\det(\mathbf{A})$
  - If  $\mathbf{B}$  results from  $\mathbf{A}$  by multiplying one row with a number  $c$ , then  $\det(\mathbf{B}) = c \cdot \det(\mathbf{A})$
  - If  $\mathbf{B}$  results from  $\mathbf{A}$  by adding a multiple of one row to another row, then  $\det(\mathbf{B}) = \det(\mathbf{A})$
- **Transpose:**  $\det(\mathbf{A}^T) = \det(\mathbf{A})$
- **Multiplication:**  $\det(\mathbf{A} \cdot \mathbf{B}) = \det(\mathbf{A}) \cdot \det(\mathbf{B})$
- Does **not** apply to addition!  $\det(\mathbf{A} + \mathbf{B}) \neq \det(\mathbf{A}) + \det(\mathbf{B})$




# Determinant: Applications

- Compute **Eigenvalues**:  
Solve the characteristic polynomial  $\det(\mathbf{A} - \lambda \cdot \mathbf{I}) = 0$
- **Area and Volume**:  $\text{area} = |\det(\mathbf{A})|$



$$\mathbf{A} = \begin{bmatrix} a & b & c \\ d & e & f \\ g & h & i \end{bmatrix}$$

( $r_i$  is  $i$ -th row)



# Orthonormal Matrix

- A matrix  $Q$  is **orthonormal** iff its column (row) vectors represent an **orthonormal** basis

$$q_{*i}^T \cdot q_{*j} = \begin{cases} 1 & \text{if } i = j \\ 0 & \text{if } i \neq j \end{cases}, \forall i, j$$

- As linear transformation, it is **norm** preserving
- Some properties:
  - The transpose is the inverse  $QQ^T = Q^TQ = I$
  - Determinant has unity norm (§ 1)

$$1 = \det(I) = \det(Q^T Q) = \det(Q)\det(Q^T) = \det(Q)^2$$

# Rotation Matrix

- A Rotation matrix is an orthonormal matrix with  $\det = +1$

- 2D Rotations  $R(\theta) = \begin{bmatrix} \cos(\theta) & -\sin(\theta) \\ \sin(\theta) & \cos(\theta) \end{bmatrix}$

- 3D Rotations along the main axes

$$R_x(\theta) = \begin{bmatrix} 1 & 0 & 0 \\ 0 & \cos(\theta) & -\sin(\theta) \\ 0 & \sin(\theta) & \cos(\theta) \end{bmatrix} \quad R_y(\theta) = \begin{bmatrix} \cos(\theta) & 0 & -\sin(\theta) \\ 0 & 1 & 0 \\ \sin(\theta) & 0 & \cos(\theta) \end{bmatrix}$$

- **IMPORTANT: Rotations are not commutative**

$$R_x\left(\frac{\pi}{4}\right) \cdot R_y\left(\frac{\pi}{4}\right) = \begin{bmatrix} 0.707 & 0 & -0.707 \\ -0.5 & 0.707 & -0.5 \\ 0.5 & 0.707 & 0.5 \end{bmatrix}, \quad R_x\left(\frac{\pi}{4}\right) \cdot R_y\left(\frac{\pi}{4}\right) \cdot \begin{bmatrix} 1 \\ 2 \\ 3 \end{bmatrix} = \begin{bmatrix} -1.414 \\ 0.586 \\ 3.414 \end{bmatrix}$$

$$R_y\left(\frac{\pi}{4}\right) \cdot R_x\left(\frac{\pi}{4}\right) = \begin{bmatrix} 0.707 & -0.5 & -0.5 \\ 0 & 0.707 & -0.707 \\ 0.707 & 0.5 & 0.5 \end{bmatrix}, \quad R_y\left(\frac{\pi}{4}\right) \cdot R_x\left(\frac{\pi}{4}\right) \cdot \begin{bmatrix} 1 \\ 2 \\ 3 \end{bmatrix} = \begin{bmatrix} -1.793 \\ 0.707 \\ 3.207 \end{bmatrix}$$

# Matrices to Represent Affine Transformations

- A general and easy way to describe a 3D transformation is via matrices

$$\mathbf{A} = \begin{pmatrix} \mathbf{R} & \mathbf{t} \\ \mathbf{0} & 1 \end{pmatrix} \quad \mathbf{A}^{-1} = \begin{pmatrix} \mathbf{R}^T & -\mathbf{R}^T \mathbf{t} \\ \mathbf{0} & 1 \end{pmatrix} \quad \mathbf{p} = \begin{pmatrix} \mathbf{t} \\ 1 \end{pmatrix}$$


Translation Vector

Rotation Matrix

- Takes naturally into account the non-commutativity of the transformations
- Homogeneous coordinates


# Combining Transformations

- A simple interpretation: chaining of transformations (represented as homogeneous matrices)
  - Matrix  $\mathbf{A}$  represents the pose of a **robot** in the space
  - Matrix  $\mathbf{B}$  represents the position of a sensor on the robot
  - The **sensor** perceives an **object** at a given location  $\mathbf{p}$ , in its own frame [the sensor has no clue on where it is in the world]
  - Where is the object in the global frame?



# Combining Transformations


- A simple interpretation: chaining of transformations (represented as homogeneous matrices)
  - Matrix  $\mathbf{A}$  represents the pose of a **robot** in the space
  - Matrix  $\mathbf{B}$  represents the position of a sensor on the robot
  - The **sensor** perceives an **object** at a given location  $\mathbf{p}$ , in its own frame [the sensor has no clue on where it is in the world]
  - Where is the object in the global frame?



$\mathbf{B}\mathbf{p}$  gives the pose of the object wrt the robot

# Combining Transformations

- A simple interpretation: chaining of transformations (represented as homogeneous matrices)
  - Matrix  $\mathbf{A}$  represents the pose of a **robot** in the space
  - Matrix  $\mathbf{B}$  represents the position of a sensor on the robot
  - The **sensor** perceives an **object** at a given location  $\mathbf{p}$ , in its own frame [the sensor has no clue on where it is in the world]
  - Where is the object in the global frame?



$\mathbf{Bp}$  gives the pose of the object wrt the robot

$\mathbf{ABp}$  gives the pose of the object wrt the world

# Positive Definite Matrix

- The analogous of positive number
- Definition  $M > 0$  iff  $z^T M z > 0 \forall z \neq 0$
- Example
  - $M_1 = \begin{bmatrix} 1 & 0 \\ 0 & 1 \end{bmatrix}$ ,  $\begin{bmatrix} z_1 & z_2 \end{bmatrix} \begin{bmatrix} 1 & 0 \\ 0 & 1 \end{bmatrix} \begin{bmatrix} z_1 \\ z_2 \end{bmatrix} = z_1^2 + z_2^2 > 0$



# Positive Definite Matrix

- Properties
  - **Invertible**, with positive definite inverse
  - All real **eigenvalues**  $> 0$
  - **Trace** is  $> 0$
  - **Cholesky** decomposition  $A = LL^T$

# Linear Systems (1)

$$\mathbf{Ax} = \mathbf{b}$$



## Interpretations:

- A set of linear equations
- A way to find the coordinates  $\mathbf{x}$  in the reference system of  $\mathbf{A}$  such that  $\mathbf{b}$  is the result of the transformation of  $\mathbf{Ax}$
- Solvable by Gaussian elimination

# Linear Systems (2)

$$\mathbf{Ax} = \mathbf{b}$$

## Notes:

- Many efficient solvers exist, e.g., conjugate gradients, sparse Cholesky decomposition
- One can obtain a reduced system  $(\mathbf{A}', \mathbf{b}')$  by considering the matrix  $(\mathbf{A}, \mathbf{b})$  and suppressing all the rows which are linearly dependent
- Let  $\mathbf{A}'\mathbf{x}=\mathbf{b}'$  the reduced system with  $\mathbf{A}':n'\times m$  and  $\mathbf{b}':n'\times 1$  and  $\text{rank } \mathbf{A}' = \min(n', m)$  rows   columns
- The system might be either over-constrained ( $n' > m$ ) or under-constrained ( $n' < m$ )

# Over-Constrained Systems

- “More (indep) equations than variables”
- An over-constrained system does not admit an **exact solution**
- However, if  $\text{rank } \mathbf{A}' = \text{cols}(\mathbf{A})$  one often computes a **minimum norm solution**

$$\mathbf{x} = \underset{\mathbf{x}}{\text{argmin}} \|\mathbf{A}'\mathbf{x} - \mathbf{b}'\|$$

Note: rank = Maximum number of linearly independent rows/columns

# Under-Constrained Systems

- “More variables than (indep) equations”
- The system is **under-constrained** if the number of linearly independent rows of  $\mathbf{A}'$  is smaller than the dimension of  $\mathbf{b}'$
- An under-constrained system admits infinite solutions
- The degree of these infinite solutions is  $cols(\mathbf{A}') - rows(\mathbf{A}')$

# Jacobian Matrix

- It is a **non-square matrix**  $n \times m$  in general
- Given a vector-valued function



$$f(\mathbf{x}) = \begin{bmatrix} f_1(\mathbf{x}) \\ f_2(\mathbf{x}) \\ \vdots \\ f_m(\mathbf{x}) \end{bmatrix}$$

- Then, the **Jacobian matrix** is defined as

$$\mathbf{F}_{\mathbf{x}} = \begin{bmatrix} \frac{\partial f_1}{\partial x_1} & \frac{\partial f_1}{\partial x_2} & \cdots & \frac{\partial f_1}{\partial x_n} \\ \frac{\partial f_2}{\partial x_1} & \frac{\partial f_2}{\partial x_2} & \cdots & \frac{\partial f_2}{\partial x_n} \\ \vdots & \vdots & \cdots & \vdots \\ \frac{\partial f_m}{\partial x_1} & \frac{\partial f_m}{\partial x_2} & \cdots & \frac{\partial f_m}{\partial x_n} \end{bmatrix}$$

# Jacobian Matrix

- It is the orientation of the **tangent plane** to the vector-valued function at a given point



- Generalizes the gradient** of a scalar valued function

# Further Reading

- A “quick and dirty” guide to matrices is the Matrix Cookbook available at:  
<http://matrixcookbook.com>